

AD-A090 430


ARMY ELECTRONICS RESEARCH AND DEVELOPMENT COMMAND WS--ETC F/6 4/1
RELATIONSHIPS BETWEEN EXTINCTION AND MASS CONTENT OF ATMOSPHERE--ETC(U)
JUN 80 R 6 PINNICK, S 6 JENNINGS

UNCLASSIFIED

NL

1 of 1
AD-A090 430

END
DATE
FILMED
11 80
DTIC



*PINNICK, JENNINGS and
CHYLEK

AD A090430

RELATIONSHIPS BETWEEN EXTINCTION AND MASS
CONTENT OF ATMOSPHERIC FOG AND MILITARY SMOKE, (U)

RONALD G. PINNICK, PhD
S. GERARD JENNINGS, PhD
ATMOSPHERIC SCIENCES LABORATORY
WHITE SANDS MISSILE RANGE, NEW MEXICO 88002

PETR CHYLEK, PhD
CENTER FOR EARTH AND PLANETARY PHYSICS, HARVARD UNIVERSITY
CAMBRIDGE, MA 02138

INTRODUCTION:

A possible relation between the extinction (or atmospheric visibility) and particle mass content of fogs and aerosols has been a subject of interest to many scientists since the beginning of the century. If such a relation exists, it would be of considerable practical interest since then remote sensing extinction measurements would yield information concerning the fog or aerosol mass content, and vice versa, from mass content measurements (which are more easily done at various meteorological stations) one could deduce the extinction properties of fogs and aerosols--at least at definite wavelengths.

All previous attempts to relate the extinction (or visibility) to the mass content of fogs and aerosols were only partially successful since all derived relations also turned out to be functions of the particle size distribution, and thus the derived relation varied from one case to another.

We will show that at definite wavelengths, which are determined primarily by the size of the largest particles present in a given polydispersion of fog or smoke aerosol particles, there exists an unambiguous relation between the extinction and mass content which is independent of the form of the particle size distribution.

80 10 17 308

*PINNICK, JENNINGS and
CHÝLEK

EXTINCTION AND MASS CONTENT OF FOG AND SMOKE AEROSOL:

Consider a polydispersion of spherical aerosol particles characterized by a size distribution $n(r)$ and refractive index m . We want to derive relationships between the aerosol extinction coefficient σ_e and the aerosol mass content M given by

$$\sigma_e = \int \pi r^2 Q_e(m, x) n(r) dr \quad (1)$$

$$M = \rho \int \frac{4}{3} \pi r^3 n(r) dr \quad (2)$$

where $Q_e(m, x)$ is the Mie efficiency factor for extinction for a particle with refractive index m and size parameter $x = 2\pi r/\lambda$ at the radiation wavelength λ and ρ is the aerosol density.

It can be seen from the integrals in equations (1) and (2) that for the extinction coefficient σ_e and mass content M to be uniquely related for arbitrary particle size distributions $n(r)$, we must be able to approximate the Mie extinction efficiency factor $Q_e(m, x)$ by a linear function of particle radius $Q_e \propto r$ [or equivalently, let us say $Q_e(x) = cx$ as suggested by Chýlek].^{1,2} Then the right-hand side of equation (1) contains the integral $\int r^3 n(r) dr$ and combining equations (1) and (2) yields

$$\sigma_e = \frac{3\pi c}{2\lambda \rho} M. \quad (3)$$

Unfortunately, the task of finding a wavelength for which the $Q_e = cx$ approximation is a good one is formidable because in general the extinction efficiency is a rather complicated function of particle size, refractive index, and wavelength. Nonetheless, examination of the form of the Mie efficiency function $Q_e(x)$ for fog droplets, phosphoric acid droplets, HC, and carbon smoke particles for wavelengths ranging from the visible through the middle infrared reveals that at particular wavelengths the $Q_e(x) = cx$ approximation is acceptable.

An example of such a wavelength for fog droplets is shown in figure 1 where the Mie extinction efficiency function at $\lambda = 11\mu m$ is well approximated by $Q_e = cx$, up to a maximum value $x_m \approx 8$

(131)

A

*PINNICK, JENNINGS and
CHÝLEK

(corresponding to a maximum radius $r_m \approx 14 \mu m$). Therefore, under the constraint that fog droplets have radii not exceeding $r_m = 14 \mu m$ (which is not unrealistic for radiation and most advection fogs), we expect a linear relation between extinction at $\lambda = 11 \mu m$ and liquid water content of fog according to equation (3).




Figure 1. The efficiency factor for extinction Q_e for water droplets versus size parameter x at a wavelength $\lambda = 11 \mu m$ (index of refraction $m = 1.153 - 0.0968i$). The efficiency factor can be approximated by a straight line $Q_e = cx$ providing $x \leq x_m$. The approximation overestimates the exact value of Q_e for some size parameters, but underestimates it for others (still with $x \leq x_m$). These two errors tend to cancel, leading to the extinction coefficient being linearly related to the liquid water content of fogs according to equation (3).

We should point out that this linear approximation of the Mie extinction efficiency is distinctly different than a Rayleigh approximation, which requires the particle size (with respect to wavelength) and phase shift be small ($|m - 1| x \ll 1$). Clearly, this Rayleigh condition is not satisfied for size parameters as large as $x = 8$ as in figure 1.

*PINNICK, JENNINGS and
CHYLEK

In the next sections we apply this approximation procedure and the resulting extinction-mass content prediction (3) to polydispersions of particles having widely different optical properties: atmospheric fog, phosphoric acid, HC, and carbon smokes. The success achieved with relation (3) attests to its simplifying impact on the gamut of radiative transfer problems in which atmospheric fog or aerosols play a role. Thus, our relation (3) connecting particulate extinction to particulate mass content should prove useful in such diverse research areas as earth climate, remote sensing, and electro-optical communications.

APPLICATION TO ATMOSPHERIC FOG:

Before we can have confidence in applying the linear relation (3) between extinction and liquid water content of fog we should test its validity with existing measurements that are available. Since simultaneous infrared extinction and liquid water content measurements in fog do not yet exist, we calculated the volume extinction coefficient σ_e using Mie theory and the liquid water content M for 341 measured size distributions of fog droplets reported in the literature.^{9, 10, 22, 13, 14, 21} The fog measurements were chosen to represent a wide range of meteorological conditions ranging from maritime and continental advection fogs^{13, 14, 9} to inland radiation fogs.^{9, 10, 22, 21}

The results of these calculations together with the derived approximation equation (3) at $\lambda = 11\mu\text{m}$ are shown in figure 2. Points showing the results of the Mie calculations are always within a factor 2 of the straight line showing the equation (3) approximation ($\sigma_e = 128W$) even though the extinction coefficient and liquid water content ranges over almost four orders of magnitude with the considered size distributions. In view of the fact that the $\sigma_e = 128W$ relation is not a result of any fit to experimental points, but rather an absolute calculation (by absolute we mean that no free fitting parameter is involved) using equation (3), we consider the agreement to be acceptable.

On the other hand, we do not expect a unique extinction-liquid content relationship at all wavelengths. To demonstrate this point, we present Mie calculations for the same 341 droplet distributions at $\lambda = 4\mu\text{m}$ in figure 3. For a given liquid water content, the extinction coefficient varies by more than an order of magnitude as a function of the size distribution.

132

*PINNICK, JENNINGS and
CHYLEK




Figure 2. Variation of particulate extinction coefficient (at $\lambda = 11 \mu m$) with liquid water content W . The points were determined from 341 size distribution measurements of atmospheric fog and haze made at different geographic locales and under a variety of meteorological conditions [open circles denote measurements of Pinnick,²¹ plus symbols of Garland and Roach^{9,10,22}, square symbols of Kumai,¹³ and solid circle symbols of Kunkel¹⁴]; the straight line labeled $\sigma_e = 128W$ was determined from equation (3).

APPLICATION TO PHOSPHORIC ACID AND RP SMOKE:

Phosphoric acid smoke differs from atmospheric fog in two important respects. First, smoke particles are much smaller than fog droplets; and second, they are generally strongly absorbing at infrared wavelengths. These two characteristics result in Mie efficiency factors Q_e being well approximated by $Q_e = cx$ for most infrared wavelengths, and consequently, we expect the extinction-mass relation (3) should be applicable throughout the infrared.

*PINNICK, JENNINGS and
CHYLEK




Figure 3. Variation of particulate extinction coefficient (at $\lambda = 4 \mu\text{m}$) with liquid water content for the 341 size distributions of fog and haze considered in figure 2.

To test this contention we have compared the ratio of the extinction coefficient to mass content σ_e/M predicted according to our size-distribution-independent relation (3) to values of σ_e/M determined from the transmission and mass content measurements of Milham.^{15*} The results, which are seen in figures 4 and 5, show

*The extinction coefficients σ_e were derived from transmission measurements for which forward scattering corrections⁷ and multiple scattering corrections should be considered. Forward scattering corrections arise from singly scattered photons that enter the detector along with the unscattered (direct) radiation due to the finite angular aperture of the detector. Similarly, multiple scattering corrections arise from signal contributed by multiply-scattered photons. Both these effects cause increased detector signal and hence result in a smaller inferred extinction coefficient if they are not considered. We estimate the forward scatter corrections for the experimental setup used by Milham¹⁵ to be not more than 3 percent and have neglected them; however, no attempt was made to make quantitative estimates of multiple scatter corrections.

134

*PINNICK, JENNINGS and
CHYLEK

agreement of predicted and measured σ_e/M values to within 30 percent throughout the $3\mu\text{m}$ to $5\mu\text{m}$ and $8\mu\text{m}$ to $12\mu\text{m}$ atmospheric window spectral regions suggesting the validity of relation (3) for phosphoric acid smoke. The extinction-mass measurements in figures 4 and 5 can only suggest (and not prove) the validity of relation (3), as the measurements are for only one particular size distribution of phosphoric acid particles.




Figure 4. Values of the ratio of aerosol extinction coefficient to aerosol mass content (σ_e/M in m^2g^{-1}) predicted according to the size-distribution-independent linear relation (3) (open circles), and measured by Milham¹⁵ (smooth curve) for 60 percent phosphoric acid aerosol. Good agreement between prediction and measurement is obtained throughout the $3\mu\text{m} \leq \lambda \leq 5\mu\text{m}$ spectral region.

In contrast to the phosphoric acid results, comparison of the relation (3) to Milham's¹⁵ measurements on RP smoke (figure 6) show relatively poor agreement in the $8\mu\text{m}$ to $12\mu\text{m}$ spectral region. The reason for the poor agreement is that the burning of RP/WP smokes apparently results in production of an unknown chemical species¹⁶ whose refractive indexes cannot be approximated by those of phosphoric acid [which was assumed in determination of $c(\lambda)$ in relation (3)].

*PINNICK, JENNINGS and
CHYLEK




Figure 5. Same as figure 4 except for the $8\mu\text{m} \leq \lambda \leq 12\mu\text{m}$ spectral region.

APPLICATION TO HC SMOKE:

Like phosphoric acid and RP smoke particles, HC smoke particles are much smaller than fog droplets. However, they are much weaker absorbers than these other smokes at most infrared wavelengths, and as a result the $Q_e = cx$ approximation is less accurate.

This inaccuracy is reflected in the comparison of σ_e/M values predicted according to equation (3) with measurements of Milham¹⁵ in figures 7 and 8. For some wavelengths in the $3\mu\text{m}$ to $5\mu\text{m}$ and $8\mu\text{m}$ to $12\mu\text{m}$ spectral regions the disagreement between prediction and measurement is as much as a factor 2.5. Thus, although we expect relation (3) to be successful throughout the infrared for strongly absorbing phosphoric acid smoke, for weakly absorbing HC smoke our relation (3) should only be applied for wavelengths around $\lambda = 4\mu\text{m}$ and $\lambda = 11\mu\text{m}$.

*PINNICK, JENNINGS and
CHÝLEK




Figure 6. Same as figure 5 except for RP smoke rather than phosphoric acid. In this case, measurements are shown for two different aerosol mass loadings (M is the aerosol mass content and L the transmission chamber path length).

APPLICATION TO CARBON SMOKE:

Caution must be exercised in applying the aerosol extinction coefficient-mass content relation (3) to carbon smoke because in general the particles are irregular in shape and the derivation of equation (3) requires the assumption of spherical homogeneous particles. We know from extinction and scattering measurements on slightly irregular particles by Greenberg,¹¹ Zerull,²⁷ and Pinnick,²⁰ that providing their equivalent size parameters are less than $x = 3$, they scatter like spheres. If carbon smoke particles were only slightly irregular and if they had size parameters less than $x = 3$, we might expect that our neglect of irregular particle effects in the application of equation (3) to these

*PINNICK, JENNINGS and
CHÝLEK

particles might be of little consequence. However, some carbon smoke particles, for example those measured by Roessler and Faxvog,^{2,3} consist of long extended chains of small ($\sim 0.05 \mu\text{m}$ diameter) spheres (figure 9). For these particles, we are probably not justified in approximating their extinction efficiency by some equivalent sphere extinction efficiency [as is done in determination of the parameter c in equation (3)].

Keeping in mind reservations concerning particle shape effects, we have proceeded to compare the extinction coefficient-mass relation (3) to transmission measurements made through carbon smokes from several different sources. We chose measurements reported by Hurley and Bailey^{1,2} on a hand-fired Lancashire boiler operating at full load, by Conner and Hodgkinson⁶ on an experimental smoke stack, by Roessler and Faxvog^{2,3} on acetylene smoke, and from the Handbook of the Society of Automotive Engineers (SAE),^{2,4} for diesel engine smoke emissions.




Figure 7. Values of the ratio of aerosol extinction coefficient to aerosol mass content (σ_e/M in $\text{m}^2 \text{g}^{-1}$) predicted according to the size-distribution-independent relation (3) (open circles), and measured by Milham^{1,5} (smooth curve) for HC smoke.

*PINNICK, JENNINGS and
CHYLEK




Figure 8. Same as figure 7 except for the $8\mu\text{m} \leq \lambda \leq 12\mu\text{m}$ spectral region. In this case, measurements are shown for two different aerosol mass loadings (M is the aerosol mass content and L the transmission chamber path length).




Figure 9. Scanning electron microscope micrograph showing carbon smoke particles collected onto a Nuclepore filter (from Roessler and Faxvog).²³

*PINNICK, JENNINGS and
CHYLEK

The results of the comparison are shown in table 1 where the values of σ_e/M predicted on the basis of equation (3) are generally within 20 percent of those determined from the measurements on various types of carbon smokes. Thus, even for quite nonspherical smoke particles, the relation (3) correctly predicts a relation between extinction and mass content, at least for visible wavelengths.

CONCLUSIONS:

We have shown that a linear size-distribution-independent relation should exist between the volume extinction coefficient and particulate mass content of atmospheric fog, phosphoric acid smoke, HC smoke, and carbonaceous smokes--but only at particular wavelengths. The suitable wavelengths are $\lambda \approx 11\mu\text{m}$ for fog, $\lambda \approx 3\mu\text{m}$ to $5\mu\text{m}$, $8\mu\text{m}$ to $12\mu\text{m}$ for phosphoric acid, $\lambda \approx 4\mu\text{m}$, $11\mu\text{m}$ for HC, and $\lambda \approx 0.55\mu\text{m}$ for carbon smoke. These relations can be used to uniquely connect the transmission across a path in a fog or smoke cloud to the path-integrated particulate mass content, providing forward scattering and multiple scattering corrections can be neglected.

TABLE 1

Comparison of measured and predicted values of the ratio of aerosol volume extinction coefficient to mass concentration σ_e/M for carbonaceous smoke particles. The measured values of σ_e were determined from visible transmissometer measurements under the assumption that forwardscatter and multiple scatter corrections to the inferred extinction coefficient are negligible. The measured values of smoke mass concentration M were determined from filter sampling. Also given are values of the smoke path length L and the transmissometer wavelength λ . The predicted values of σ_e/M were obtained from the size-distribution-independent relation (3) under the assumption of a particle density $\rho = 2 \text{ g cm}^{-3}$.

Reference	Mass Concentration $M(\text{g m}^{-3})$	Path Length $L(\text{m})$	Wavelength $\lambda(\mu\text{m})$	σ_e/M $(\text{m}^2\text{g}^{-1})$	Predicted [(from equation (3)] $\sigma_e/M = 3\pi c/2\lambda\rho$ $(\text{m}^2\text{g}^{-1})$
Black Smoke	0.29	0.305	0.55	8.04	9.5
(coal stoker)	1.05	0.305	0.55	8.38	
Hurley and Bailey (1958)	1.40	0.305	0.55	9.0	
Black Smoke	0.16	0.10	0.56	8.78	9.3
(oil fired smoke stack)	0.55	0.10	0.56	9.61	
Conner and Hodkinson (1967)	1.0	0.10	0.56	9.52	
Acetylene smoke	0.0155	1.835	0.514	9.93	10.2
Roessler and Faxvog (1979)	0.0527	0.10	0.514	9.96	
	0.120	1.835	0.514	9.68	
	0.165	0.10	0.514	9.60	
	0.321	0.10	0.514	10.1	
Diesel Engine Emission SAE handbook (1977)	1.0	0.10	0.55	6.79	9.5

*PINNICK, JENNINGS and
CHÝLEK

REFERENCES

1. Barteneva, O. D., and E. A. Polyakova, 1965, "A Study of Attenuation and Scattering of Light in a Natural Fog Due to its Microphysical Properties," Izv Atmospheric Oceanic Phys, 1:193-207.
2. Box, M. A., and B. H. J. McKellar, 1978, "Direct Evaluation of Aerosol Mass Loadings from Multispectral Extinction Data," Quart J Roy Meteorol Soc, 104:755-781.
3. Charlson, R. J., N. C. Ahlquist, and H. Horvath, 1968, "On the Generality of Correlation of Atmospheric Mass Concentration and Light Scatter," Atmospheric Environ, 2:455-464.
4. Chýlek, Petr, 1978, "Extinction and Liquid Water Content of Fogs," J Atmospheric Sci, 35:296-300.
5. Chýlek, Petr, J. T. Kiehl, and M. K. W. Ko, 1979, "Infrared Extinction and the Mass Concentration of Atmospheric Aerosols," Atmospheric Environ, 13:169-173.
6. Conner, W. D., and J. R. Hodgkinson, 1967, "Optical Properties and Visual Effects of Smoke-Stack Plumes," US Department of Health, Education, and Welfare, Public Health Service Publication, 999-AP-30.
7. Deepak, A., and M. A. Box, 1978, "Forwardscattering Corrections for Optical Extinction Measurements in Aerosol Media. 1: Monodispersions," Appl Opt, 17:2900-2908.
8. Eldridge, R. G., 1971, "The Relationship Between Visibility and Liquid Water Content in Fog," J Atmospheric Sci, 1183-1186.
9. Garland, J. A., 1971, "Some Fog Droplet Size Distributions Obtained by an Impaction Method," Quart J Roy Meteorol Soc, 97:483-494.
10. Garland, J. R., and I. C. Cox, 1973, "A Study of the Contribution of Pollution to Visibility in a Radiation Fog," Atmospheric Environ, 7:1079-1092.
11. Greenberg, J. M., 1972, "Absorption and Emission of Radiation by Nonspherical Particles," J Coll and Inter Sci, 39:513-519.
12. Hurley, T. F., and P. L. Bailey, 1958, "The Correlation of Optical Density with the Concentration and Composition of the Smoke Emitted from a Lancashire Boiler," J Inst Fuel, 31:534-540.
13. Kumai, M., 1973, "Arctic Fog Droplet Size Distribution and Its Effect on Light Attenuation," J Atmospheric Sci, 30:635-643.
14. Kunkel, B. A., 1971, "Fog Drop-Size Distributions Measured with a Laser Hologram Camera," J Appl Meteorol, 10:482-486.
15. Milham, M. E., 1976, "A Catalog of Optical Extinction Data for Various Aerosols/Smokes," Report ED-SP-77002, Edgewood Arsenal, Aberdeen Proving Ground, MD.

*PINNICK, JENNINGS and
CHÝLEK

16. Milham, M. E., D. H. Anderson, R. H. Frickel, and T. L. Tarnove, 1977, "New Findings on the Nature of WP/RP Smokes," Technical Report ARCSL-TR-77067, US Army ARADCOM, Chemical Systems Laboratory, Aberdeen Proving Ground, MD.
17. Noll, K. E., P. K. Mueller, and M. Imada, 1968, "Visibility and Aerosol Concentration in Urban Air," Atmospheric Environ, 2:465-475.
18. Patterson, E. M., and D. A. Gillette, 1977, "Measurements of Visibility vs Mass-Concentration for Airborne Soil Particles," Atmospheric Environ, 11:193-196.
19. Pilat, M. J., and D. S. Ensor, 1970, "Plume Opacity and Particulate Mass Concentration," Atmospheric Environ, 4:163-173.
20. Pinnick, R. G., D. E. Carroll, and D. J. Hoffman, 1976, "Polarized Light-Scattered from Monodisperse Randomly Oriented Nonspherical Aerosol Particles: Measurements," Applied Optics, 15:384-393.
21. Pinnick, R. G., D. L. Hoihjelle, G. Fernandez, E. B. Sternmark, J. D. Lindberg, S. G. Jennings, and G. B. Hoidale, 1978, "Vertical Structure in Atmospheric Fog and Haze and Its Effects on IR and Visible Extinction," J Atmospheric Sci, 35:2020-2032.
22. Roach, W. T., R. Brown, S. J. Caughly, J. A. Garland, and C. J. Readings, 1976, "The Physics of Radiation Fog: I - A Field Study," Quart J Roy Meteorol Soc, 102:313-333.
23. Roessler, D. M., and F. R. Faxvog, 1979, "Optical Properties of Acetylene Smoke at 0.5145 μ m and 10.6 μ m Wavelengths," submitted for publication in J Opt Soc Am.
24. SAE Handbook, 1977, "Diesel Engine Smoke Measurement (Steady State)," SAE J255, SAE Information Report, 871-877.
25. Tomasi, C., and F. Tampieri, 1976, "Features of the Proportionality Coefficient in the Relationship Between Visibility and Liquid Water Content in Haze and Fog," Atmosphere, 14:61-76.
26. Trabert W., 1901, "Die Extinction des Lichtes in einem truben Medium," Meteorol Z, 18:518-525.
27. Zerull, R. H., 1976, "Scattering Measurements of Dielectric and Absorbing Nonspherical Particles," Beitr Phys Atmosphere, 49:168-188.

**DA
FILM**